

Breast Diseases

EDITOR-IN-CHIEF

S. Eva Singletary, M.D.


EDITORS:

Charles M. Balch, M.D.


Gerald D. Dodd, Jr., M.D.

I. Craig Henderson, M.D.

Carlos A. Perez, M.D.



Overall survival according to estrogen receptor (ER) status. Closed squares indicate ER-negative; open squares, ER-positive (see Abstract 2-24).



Survival after a first metastasis according to estrogen receptor (ER) status. Closed squares indicate ER-negative; open squares, ER-positive (see Abstract 2-24).

- *National Cancer Data Base Report on Age and Outcome*
- *LCIS Results from NSABP Protocol B-17*
- *Pamidronate for Palliation of Bone Metastases*
- *Survival by Estrogen Receptor Status*
- *Breast-conserving Therapy After Augmentation Mammoplasty*
- *Necessity of Axillary Node Dissection*
- *Survival Impact of Radiation*
- *Legislative Mandates for Breast-conserving Surgery*

PUBLISHER'S STAFF

Vice President, Continuity Publishing
Kenneth H. Killion

Director, Editorial Development
Gretchen C. Murphy

Acquisitions Editor
Gina Byrd

**Illustrations and Permissions
Coordinator**
Lois M. Ruebensam

Developmental Editor
Catherine Flanagan

Project Supervisor, Editing
Rebecca Nordbrock

Assistant Project Supervisor
Laura M. Higgins

Freelance Staff Supervisor
Barbara M. Kelly

Director, Literature Services
Edith Podrazik, B.S.N., R.N.

Information Specialist
Terri Santo, R.N.

Information Specialist
Margery Marble, B.S., R.N.

Circulation Manager
Lynn D. Stevenson

Breast Diseases: A Year Book® Quarterly is your guide and distillation of the most important literature published worldwide. To make this quarterly, Mosby-Year Book, Inc. subscribes to and surveys nearly 950 U.S. and foreign medical and allied health journals. From these journals, the physician editors select the most important, timely, and controversial articles. Essential points are captured in 300- to 400-word abstracts, and the physician editors lend their expertise and perspective by commenting on the significance of each article.

Breast Diseases: A Year Book® Quarterly has been designed to maximize readability and to help the reader locate specific information quickly. The format features:

- Complete reference citation
- Abstracted text of the original article
- Selected illustrations and tables from the original article
- Editors' comments following each abstract
- Mini-editorials from the editors
- Historical vignettes
- Annotated bibliographies

We encourage you to place your *Breast Diseases: A Year Book® Quarterly* on the top of your stack of journals in your office. You may not have time to read all of your journals, but certainly read *Breast Diseases*. It will point out the articles you should seek in that ever-growing stack of literature.

Note: *Breast Diseases: A Year Book® Quarterly* is a literature survey service providing abstracts of articles published in the professional literature. Every effort is made to ensure the accuracy of the information presented in these pages. Neither the editors nor the publisher of *Breast Diseases: A Year Book® Quarterly* can be responsible for errors in the original materials. The editors' comments are their own opinions. Mention of specific products within this publication does not constitute endorsement.

To facilitate the use of *Breast Diseases: A Year Book® Quarterly* as a reference tool, all illustrations and tables included in this publication are now identified as they appear in the original article. This change is meant to help the reader recognize that any illustration or table appearing in *Breast Diseases* may be only one of many in the article. For this reason, figure and table numbers will often appear to be out of sequence within *Breast Diseases*.

© June 1997 by
MOSBY-YEAR BOOK, INC.

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means—electronic, mechanical, photocopying, recording, or otherwise—without prior written permission from the publisher.

Subscription Information

Vol. 8, No. 2 (July–September) 1997. *Breast Diseases: A Year Book® Quarterly* (ISSN 1043-321X) is published quarterly by Mosby-Year Book, Inc., 161 N. Clark St., Chicago, IL 60601. E-mail address: series.editorial@mosby.com Periodicals paid at St. Louis, MO, and additional mailing offices. **Postmaster:** Send address changes to *Breast Diseases: A Year Book® Quarterly*, Mosby-Year Book, Inc., 11830 Westline Industrial Drive, St. Louis, MO 63146-3318. Annual subscription rates for 1997: \$75.00 for individuals and \$112.50 for institutions. E-mail address: customer.support@mosby.com

induced neuropathy is also rare because most radiotherapists do not irradiate the axilla, particularly if the patient has had an axillary dissection. Third, pain caused by a stretching of the motor nerves in the chest, such as the thoracodorsal and pectoral nerves, is also rare in my experience. Fourth,

problems with frozen shoulders and inability to elevate the arm are markedly decreased because axillary dissections are not as extensive as they were before and, in addition, the pectoralis major muscle is saved in every mastectomy. Finally, breast reduction patients are actually some of the most

appreciative and satisfied of all patients with breast problems because of the severe discomfort they have pre-operatively and the relief experienced by the great majority of patients who undergo a reduction mammoplasty.

L.Q. Vasconez, M.D.

2-60

Reduction Mammoplasty With the Inferior Pedicle Technique: Early and Late Complications in 371 Patients

Mandrekas AD, Zambacos GJ, Anastasopoulos A, et al (Plastic Surgery Centre of Athens, Greece)

Br J Plast Surg 48:442-446, 1996

Background.—The inferior pedicle technique is one of the most popular reduction mammoplasty approaches. Patient satisfaction is high for this surgical technique. A total of 371 patients who underwent bilateral reduction mammoplasty with the inferior pedicle technique during a 10-year period were reviewed.

Methods.—Two patients had undergone a previous breast reduction. Liposuction of the breast was used in 24 selected patients as an adjunct at the beginning of surgery to make the breasts smaller and to decrease lateral breast fullness. The average patient age was 33.1 years (range, 15-67 years). The average distance from sternal notch to nipple was 30.4 cm (range, 21-43 cm). All patients received perioperative antibiotics and had suction drains until hospital discharge. Length of hospital stay was 12-24 hours.

Results.—The mean amount of breast tissue resected was 870 g per breast (range, 250-1,960 g). The mean liposuction volume removed was 160 cc (range, 100-400 cc). The amount of blood in the suction drains never exceeded 50 mL, and no blood transfu-

sions were required. Mean operative time was 3.1 hours. The overall complication rate in this cohort was 11.4% (Table 1). The pathology examination revealed breast carcinoma in 1 breast each of 2 women (0.5%) aged 45 and 54 years. Both women were referred to a breast surgeon for evaluation. Of 18 women who gave birth after surgery, 13 (72%) were able to breast-feed with decreased capacity. The other 6 patients had either milk insufficiency or masto-

dynia; their lactation was suppressed pharmacologically.

Conclusions.—Reduction mammoplasty with the inferior pedicle technique is a versatile method applicable to patients with a wide range of breast sizes. Breast contour, volume, nipple projection, nipple sensation, scar appearance, and the potential for lactation are comparable to or better than the outcome of other techniques, although the scar is longer. The complication rate

TABLE 1.—Results

	Patients
Early complications	21 (5.7%)
Haematoma	1 (0.3%)
Nipple and/or pedicle necrosis	3 (0.8%)
Wound dehiscence	17 (4.5%)
Late complications	21 (5.7%)
Fat necrosis	3 (0.8%)
Carcinoma	2 (0.5%)
Loss of sensitivity and erectility of the nipple	2 + 3* (0.5 + 0.8%)
Hypertrophic scars	12 (3.3%)
Dermoid cyst	1 (0.3%)
Marked fullness requiring secondary surgery	1 (0.3%)
Total complications	42 (11.4%)
No complications	329 (88.6%)
Total patients	371

Note: A total of 42 patients experienced 42 complications. None of the patients had more than 1 complication.

*The three patients with nipple necrosis.

(Courtesy of Mandrekas AD, Zambacos GJ, Anastasopoulos A, et al: Reduction mammoplasty with the inferior pedicle technique: Early and late complications in 371 patients. *Br J Plast Surg* 49:442-446, 1996.)

is low, and nipple sensation and lactation potential are usually maintained.

This well-written review of 371 patients who underwent inferior pedicle resection for breast reduction clearly outlines the advantages and disadvantages of this popular breast reduction technique. Since its description in 1977, the inferior pedicle technique has been well accepted among plastic surgeons as a safe and versatile method for reducing the breast. Although it is a tech-

nique that is relatively easy to teach and learn, it requires significant experience to perfect.

Other more recently described breast reduction techniques emphasize minimal scarring, but my impression is that patients are much more concerned with the overall correction of size, shape, and symmetry of the breasts. Surgeons are often more concerned with the scarring after breast reduction surgery than their patients seem to be. Furthermore, techniques such as the

inferior pedicle technique, which leaves an "inverted T" scar at the base of the breast, can be designed and executed so that no portion of the horizontal limb of the incision is exposed at either the medial or lateral aspects of the inframammary fold.

Regardless of the technique used, the surgeon should choose 1 or 2 breast reduction techniques that are safe and reliable and develop a proficiency with them.

D.P. Goldberg, M.D.

2-61

The Tuberous Breast Deformity: Classification and Treatment

von Heimburg D, Exner K, Kruft S, et al
(St Markus Hosp, Frankfurt/Main, Germany;
Johann Wolfgang Goethe Univ Teaching
Hosp, Frankfurt/Main, Germany)

Br J Plast Surg 49:339-345, 1998

Background.—Several reports have offered descriptive terminology for the so-called tuberous breast deformity, and a number of different operative procedures have been proposed. There is no consensus on nomenclature for this deformity. The degrees of tuberous breast deformity and the results of operative correction were evaluated retrospectively.

Methods.—Forty patients with 68 tuberous breast deformities were operated on at 1 institution during a 20-year period. On the basis of preoperative photographs, the breasts were classified into 4 categories: type I, hypoplasia of the lower medial quadrant; type II, hypoplasia of the lower medial and lateral quadrants with sufficient skin in the subareolar region; type III, hypoplasia of the lower medial and lateral quadrants with deficient skin in the subareolar region; and type IV, severe breast constriction with a minimal breast base (Fig 1). At least 2 years' follow-up was available in 51 breasts of 31 patients.

Results.—In the study sample, 28% of breasts had type I deformities, 26% had type II deformities, 18% had type

III deformities, and 28% had type IV deformities. Just 44% of breasts studied had areolar prolapse, which is typically




FIGURE 1.—Classification of tuberous breast deformities. (Courtesy of von Heimburg D, Exner K, Kruft S, et al: The tuberous breast deformity: Classification and treatment. *Br J Plast Surg* 49:339-345, 1998.)